arXiv:math-ph/0404051v1 23 Apr 2004

FUNDAMENTAL SOLUTIONS OF PSEUDO-DIFFERENTIAL OPERATORS OVER $p-$ADIC FIELDS

W. A. ZUNIGA-GALINDO

ABSTRACT. We show the existence of fundamental solutions for $p-$adic pseudo-differential operators with polynomial symbols.

1. INTRODUCTION

Let K be a $p-$adic field, i.e. a finite extension of $\mathbb{Q}_p$ the field of $p-$adic numbers. Let R_K be the valuation ring of K, P_K the maximal ideal of R_K, and $\overline{K}=R_K/P_K$ the residue field of K. The cardinality of $\overline{K}$ is denoted by q. For $z\in K$, $v(z)\in\mathbb{Z}\cup\{+\infty\}$ denotes the valuation of z, $|z|_K=q^{-v(z)}$ and $ac(z)=z\pi^{-v(z)}$ where π is a fixed uniformizing parameter for R_K. Let Ψ denote an additive character of K trivial on R_K but not on P_K^{-1}.

A function $\Phi:K^n\to\mathbb{C}$ is called a Schwartz-Bruhat function if it is locally constant with compact support. We denote by $\mathcal{S}(K^n)$ the $\mathbb{C}$-vector space of Schwartz-Bruhat functions over K^n.The dual space $\mathcal{S}'(K^n)$ is the space of distributions over K^n. Let $f=f(x)\in K[x]$, $x=(x_1,..,x_n)$, be a non-zero polynomial, and β a complex number satisfying $\mathrm{Re}(\beta)>0$. If $x=(x_1,...,x_n)$, $y=(y_1,...,y_n)\in K^n$, we set $[x,y]:=\sum_{i=1}^n x_iy_i$.

A p-adic pseudo-differential operator $f(\partial,\beta)$, with symbol $|f|_K^\beta$, is an operator of the form

$$\begin{array}{rccc} f(\partial,\beta): & \mathcal{S}(K^n) & \to & \mathcal{S}(K^n) \\ & \Phi & \to & \mathcal{F}^{-1}\left(|f|_K^\beta\,\mathcal{F}(\Phi)\right), \end{array} \tag{1.1}$$

where

$$\begin{array}{rccc} \mathcal{F}: & \mathcal{S}(K^n) & \to & \mathcal{S}(K^n) \\ & \Phi & \to & \int\limits_{K^n}\Psi(-[x,y])\,\Phi(x)\,dx \end{array} \tag{1.2}$$

is the Fourier transform. The operator $f(\partial,\beta)$ has self-adjoint extension with dense domain in $L^2(K^n)$. We associate to $f(\partial,\beta)$ the following p-adic pseudo-differential equation:

$$f(\partial,\beta)u=g,\ \ g\in\mathcal{S}(K^n). \tag{1.3}$$

A fundamental solution for (1.3) is a distribution E such that $u=E*g$ is a solution.

The main result of this paper is the following.

1991 *Mathematics Subject Classification.* Primary 46S10, 11S40.

Key words and phrases. non-archimedean functional analysis, pseudo-differential operators, Igusa's local zeta function.

Theorem 1.1. *Every p-adic pseudo-differential equation* $f(\partial,\beta)u=g$, *with* $f(x)\in K[x_1,..,x_n]\setminus K$, $g\in\mathcal{S}(K^n)$, *and* $\beta\in\mathbb{C}$, $\operatorname{Re}(\beta)>0$, *has a fundamental solution* $E\in\mathcal{S}'(K^n)$.

The $p-$adic pseudo-differential operators occur naturally in $p-$adic quantum field theory [11], [6]. Vladimirov showed the existence of a fundamental solutions for symbols of the form $|\xi|_K^{\alpha}$, $\alpha>0$ [10], [11]. In [7], [6] Kochubei showed explicitly the existence of fundamental solutions for operators with symbols of the form $|f(\xi_1,..,\xi_n)|_K^{\alpha}$, $\alpha>0$, where $f(\xi_1,..,\xi_n)$ is a quadratic form satisfying $f(\xi_1,..,\xi_n)\neq 0$ if $|\xi_1|_K+..+|\xi_n|_K\neq 0$. In [8] Khrennikov considered spaces of functions and distributions defined outside the singularities of a symbol, in this situation he showed the existence of a fundamental solution for a $p-$adic pseudo-differential equation with symbol $|f|_K\neq 0$. The main result of this paper shows the existence of fundamental solutions for operators with polynomial symbols. Our proof is based on a solution of the division problem for $p-$adic distributions. This problem is solved by adapting the ideas developed by Atiyah for the archimedean case [1], and Igusa's theorem on the meromorphic continuation of local zeta functions [3], [4]. The connection between local zeta functions (also called Igusa's local zeta functions) and fundamental solutions of $p-$adic pseudo-differential operators has been explicitly showed in particular cases by Jang and Sato [5], [9]. In [9] Sato studies the asymptotics of the Green function G of the following pseudo-differential equation

$$\left(f(\partial,1)+m^2\right)u=g,\ \ m>0. \tag{1.4}$$

The main result in [9, theorem 2.3] describes the asymptotics of $G(x)$ when the polynomial f is a relative invariant of some prehomogeneous vector spaces (see e.g. [3, Chapter 6]). The key step is to establish a connection between the Green function $G(x)$ and the local zeta function attached to f.

All the above mentioned results suggest a deep connection between Igusa's work on local zeta functions (see e.g. [3]) and $p-$adic pseudo-differential equations.

2. Local zeta functions and division of distributions

The local zeta function associated to f is the distribution

$$\langle |f|_K^s,\Phi\rangle=\int\limits_{K^n\setminus f^{-1}(0)}\Phi(x)\,|f(x)|_K^s\,dx, \tag{2.1}$$

where $\Phi\in\mathcal{S}(K^n)$, $s\in\mathbb{C}$, $\operatorname{Re}(s)>0$, and dx is the Haar of K^n normalized so that $vol(R_K^n)=1$. The local zeta functions were introduced by Weil [12] and their basic properties for general f were first studied by Igusa [3], [4]. A central result in the theory of local zeta functions is the following.

Theorem 2.1 (Igusa, [3, Theorem 8.2.1])**.** *The distribution* $|f|_K^s$ *admits a meromorphic continuation to the complex plane such that* $\langle |f|_K^s,\Phi\rangle$ *is a rational function of* q^{-s} *for each* $\Phi\in\mathcal{S}(K^n)$. *In addition the real parts of the poles of* $|f|_K^s$ *are negative rational numbers.*

The archimedean counterpart of the previous theorem was obtained jointly by Bernstein and Gelfand [2], independently by Atiyah [1]. The following lemma is a consequence of the previous theorem.

Lemma 2.1. *Let $f(x) \in K[x_1,..,x_n]$ be a non-constant polynomial, and β a complex number satisfying $\mathrm{Re}(\beta) > 0$. Then there exists a distribution $T \in \mathcal{S}'(K^n)$ satisfying $|f|_K^\beta T = 1$.*

Proof. By theorem 2.1 $|f|_K^s$ has a meromorphic continuation to $\mathbb{C}$ such that $\langle |f|_K^s, \Phi\rangle$ is a rational function of q^{-s} for each $\Phi \in \mathcal{S}(K^n)$. Let

$$|f|_K^s = \sum_{m\in\mathbb{Z}} c_m (s+\beta)^m \tag{2.2}$$

be the Laurent expansion at $-\beta$ with $c_m \in \mathcal{S}'(K^n)$ for all m. Since the real parts of the poles of $|f|_K^s$ are negative rational numbers by theorem 2.1, it holds that $|f|_K^{s+\beta} = |f|_K^\beta \ |f|_K^s$ is holomorphic at $s = -\beta$. Therefore $|f|_K^\beta c_m = 0$ for all $m < 0$ and

$$|f|_K^{s+\beta} = c_0 |f|_K^\beta + \sum_{m=1}^{\infty} c_m |f|_K^\beta (s+\beta)^m . \tag{2.3}$$

By using the Lebesgue lemma and (2.3) it holds that

$$\begin{aligned} \lim_{s\to-\beta} \left\langle |f|_K^{s+\beta}, \Phi \right\rangle &= \int_{K^n} \Phi(x)\,dx = \langle 1, \Phi\rangle \\ &= c_0 |f|_K^\beta . \end{aligned} \tag{2.4}$$

Therefore we can take $T = c_0$. ∎

If $T \in \mathcal{S}'(K^n)$ we denote by $\mathcal{F}T \in \mathcal{S}'(K^n)$ the Fourier transform of the distribution S, i.e. $\langle \mathcal{F}T, \Phi\rangle = \langle S, \mathcal{F}(\Phi)\rangle$, $\Phi \in \mathcal{S}(K^n)$.

3. Proof of the main result

By lemma 2.1 there exists a $T \in \mathcal{S}'(K^n)$ such that $|f|_K^\beta T = 1$. We set $E = \mathcal{F}^{-1}T \in \mathcal{S}'(K^n)$ and assert that E is a fundamental solution for (1.3). This last statement is equivalent to assert that $\mathcal{F}(\Phi) = (\mathcal{F}E)\mathcal{F}(g)$ satisfies $|f|_K^\beta \mathcal{F}(\Phi) = \mathcal{F}(g)$. Since $|f|_K^\beta \mathcal{F}(\Phi) = |f|_K^\beta (\mathcal{F}E)\mathcal{F}(g) = |f|_K^\beta T\, \mathcal{F}(g) = \mathcal{F}(g)$, we have that E is a fundamental solution for (1.3).

4. Operators with twisted symbols

Let $\chi : R_K^\times \to \mathbb{C}$ be a non-trivial multiplicative character, i.e. a homomorphism with finite image, where $R_K^\times$ is the group of units of R_K. We put formally $\chi(0) = 0$. If $f(x) \in K[x_1,..,x_n] \setminus K$, we say that $\chi(ac(f))|f|_K^\beta$, with $\beta \in \mathbb{C}$, $\mathrm{Re}(\beta) > 0$, is a *twisted symbol*, and call the pseudo-differential operator

$$\Phi \to f(\partial,\beta,\chi)\Phi = \mathcal{F}^{-1}\left(\chi(ac(f))|f|_K^\beta \mathcal{F}(\Phi)\right), \Phi \in \mathcal{S}(K^n), \tag{4.1}$$

a *twisted operator*. Since the distribution $\chi(ac(f))|f|_K^\beta$ satisfies all the properties stated in theorem 2.1 (cf. [3, Theorem 8.2.1]), theorem 1.1 generalizes literally to the case of twisted operators. In [6, chapter 2] Kochubei showed explicitly the existence of fundamental solutions for twisted operators in some particular cases.

References

[1] M. F. Atiyah, Resolution of singularities and division of distributions, Comm. Pure Appl. Math. 23 (1970), 145-150.

[2] I. N. Bernstein and S. I. Gelfand, Meromorphic property of the functions P^{λ}, Functional Anal. Appl., 3 (1969), 68-69.

[3] Igusa Jun-Ichi, An introduction to the theory of local zeta functions, AMS /IP studies in advanced mathematics, v. 14, 2000.

[4] Igusa J., Complex powers and asymptotic expansions, I Crelles J. Math., 268/269 (1974), 110-130; II, ibid., 278/279 (1975), 357-368.

[5] Y. Jang, An asymptotic expansion of the $p-$adic Green function, Tohoku Math. J., 50 (1998), 229-242.

[6] A. N. Kochubei, Pseudodifferential equations and stochastics over non-archimedean fields, Marcel Dekker, 2001.

[7] A. N. Kochubei, Fundamental solutions of pseudo-differential equations associated with $p-$adic quadratic forms, Izvestiya Math., 62 (1998), 1169-1188.

[8] A. Khrennikov, Fundamental solutions over the field of p-adic numbers, St. Peterburgh Math. J., 4 (1993), 613-628.

[9] Fumihiro Sato, $p-$adic Green functions and zeta functions, Commentarii Mathematici Universitatis Sancti Pauli, 51 (2002), 79-97.

[10] V. S. Vladimirov, On the spectrum of some pseudo-differential operators over $p-$adic number field, Algebra and Analysis, 2, 107-124 (1990).

[11] V. S. Vladimirov, I. V. Volovich and E. I. Zelenov, $p-$adic Analysis and mathematical physics, World Scientific, Singapore, 1994.

[12] A. Weil, Sur la formule de Siegel dans le théorie des groupes classiques, Acta Math. 113 (1965), 1-87.

Department of Mathematics and Computer Science, Barry University, 11300 N.E. Second Avenue, Miami Shores, Florida 33161, USA

E-mail address: wzuniga@mail.barry.edu